


Halogen bonding sensors for environmental pollutants

Oxford researchers have developed a novel pillararene host molecule for the optical detection of dicarboxylates and other environmentally harmful substances.

Applications: Environmental sensing, molecular Recognition, chemical purification



Features	Benefits
Enhanced sensing	Improved detection
Adjustable componentry to translate the sensing properties to a readable or quantifiable output	Integrated halogen bonding recognition sites with bright fluorophores, meaning a sensor can identify pollutants through distinct fluorescence response
Detection indicator can be switch-on or switch-off, as required	Superior binding affinity compared to traditional hydrogen-bonding alternatives
	This ensures that a user gets a clear signal when target compounds are detected

Features	Benefits
	Particularly effective in aqueous-organic media, where the hydrophobic effect significantly enhances the receptor's sensitivity
	Allows the end user to dictate the preferred mechanism
	Demonstrates a powerful strategy for sensing both charged and neutral environmental contaminants

Patented and Available for

- Co-development
- Consulting
- Licensing

To find out more, contact enquiries@innovation.ox.ac.uk and quote the project number.

The above information is provided 'as is' without conditions or warranties. Oxford University Innovation makes no representation and gives no warranty that it is the owner of the intellectual property rights in the technology described. © Oxford University Innovation Ltd 2026, Buxton Court, 3 West Way, Oxford, OX2 0JB.