


Synthesising agarwood aroma compounds through selective oxidation of alpha-bulnesene

Creating the scent of agarwood by selectively changing the alpha-bulnesene compound.

Applications: Fragrance, Cosmetics



Features

Biotransformation of α -bulnesene by cytochrome P450BM3 into key aroma compounds of agarwood. The starting material is derived from patchouli oil

Formation of the lactone involves a one-pot enzyme cascade using two variants of the same enzyme (cytochrome P450BM3)

Engineered P450BM3 variants with high activity and regioselectivity (>90%) for C15 oxidation of α -bulnesene

Benefits

Agarwood-producing Aquilaria trees have been listed as threatened species since 2006. This process provides a more sustainable source for agarwood aroma compounds by using patchouli, a much more abundant resource

More robust and low-cost process compared to conventional cascades that use different enzymes

Much higher yield compared to the resource intensity of conventional extraction, where typically 70kg of agarwood yields just 20g of oil

Features

Synthesis of both the aldehyde and lactone by the oxidation of α -bulnesene

Benefits

The hydroxylated products of α -bulnesene could additionally have wider flavour, fragrance or biological properties

Available for

- Co-development
- Consulting
- Licensing

To find out more, contact enquiries@innovation.ox.ac.uk and quote the project number.

The above information is provided 'as is' without conditions or warranties. Oxford University Innovation makes no representation and gives no warranty that it is the owner of the intellectual property rights in the technology described. © Oxford University Innovation Ltd 2026, Buxton Court, 3 West Way, Oxford, OX2 0JB.